

# ELECTROCHEMISTRY

## Electrochemistry

electrons) are required. Hence,  $3 \times 2.00 = 6.00$  F are required to produce 53.96 g of Al.

(c)  $1F = 96500 \text{ C per mole e}^-$

$$\text{hence } 6.00F = 6.00 \times 96500 \text{ C}$$

$$(d) 1\text{C} = 1 \text{ amp-s (OR Q = It)}$$

$$(e) \therefore t = Q/I = 5.790 \times 10^5 \text{ C} / 10\text{A} = 57900 \text{ s} = 16.08 \text{ h}$$

$5.790 \times 10^5 \text{ C}$  are required to produce 53.96g Al

$$2 \text{ min} = 2 \times 60 = 120\text{s}$$

$$Q = It \therefore I = Q/t = \frac{5.790 \times 10^5 \text{ C}}{120 \text{ s}} (\text{A})$$

$$I = 4,825 \text{ A}$$


In general, electrolysis of aqueous solutions containing any metal ion that is a weaker oxidizing agent than  $\text{H}^+$  (or  $\text{H}_3\text{O}^+$ ) (like alkali metal and alkaline earth cations,  $\text{Al}^{3+}$ ,  $\text{Mn}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Cr}^{3+}$ ,  $\text{Fe}^{2+}$ ,  $\text{CO}^{2+}$ ,  $\text{Ni}^{2+}$ ,  $\text{Sn}^{2+}$ ,  $\text{Pb}^{2+}$ ,  $\text{Fe}^{3+}$ ) will produce  $\text{H}_2(\text{g})$  at the cathode. In other words, no reducing agent stronger than  $\text{H}_2$  is liberated during electrolysis of aqueous solutions.

## QUESTIONS

6.1 Which of the following redox reactions will be spontaneous?

- |  |                              |
|--|------------------------------|
| (a) $\text{H}_2(\text{g}) + \text{Hg}^{2+}(\text{aq}) \rightarrow 2\text{H}^+(\text{aq}) + \text{Hg}(\text{l})$          | $E^\circ = 0.851 \text{ V}$  |
| (b) $\text{Zn}(\text{s}) + 2\text{Ti}^+(\text{aq}) \rightarrow \text{Zn}^{2+}(\text{aq}) + 2\text{Ti}(\text{s})$         | $E^\circ = 0.427 \text{ V}$  |
| (c) $2\text{Al}(\text{s}) + 3\text{Ni}^{2+}(\text{aq}) \rightarrow 2\text{Al}^{3+}(\text{aq}) + 3\text{Ni}(\text{s})$    | $E^\circ = 1.43 \text{ V}$   |
| (d) $2\text{Fe}^{2+}(\text{aq}) + \text{I}_2(\text{aq}) \rightarrow 2\text{Fe}^{3+}(\text{aq}) + 2\text{I}^-(\text{aq})$ | $E^\circ = -0.235 \text{ V}$ |
| (e) $\text{Fe}(\text{s}) + \text{Cr}^{3+}(\text{aq}) \rightarrow \text{Fe}^{3+}(\text{aq}) + \text{Cr}(\text{s})$        | $E^\circ = -0.70 \text{ V}$  |
| (f) $6\text{H}^+(\text{aq}) + 2\text{Sc}(\text{s}) \rightarrow 3\text{H}_2(\text{g}) + 2\text{Sc}^{3+}$                  | $E^\circ = 2.08 \text{ V}$   |

6.2 Given the following data:



- (a) Determine the standard cell potential (for a spontaneous reaction)

- (b) Write the spontaneous cell reaction  
 (c) Write the cell diagram
- 6.3 Given the following data:



$$E^\circ = 0.00 \text{ V}$$



$$E^\circ = 0.34 \text{ V}$$



$$E^\circ = 0.96 \text{ V}$$

Explain why copper metal does not dissolve in a typical strong acid like HCl, but will dissolve in 1.0M HNO<sub>3</sub>.

- 6.4 The standard cell potential for the following reaction is 1.105V:







- (a) Write equations for the anode, cathode and overall cell reactions

- (b) Calculate  $E^\circ_{\text{Cu}^{2+}/\text{Cu}}$ , given that  $E^\circ_{\text{Zn}^{2+}/\text{Zn}} E^\circ_{\text{Cu}^{2+}/\text{Cu}} = -0.763 \text{ V}$



- 6.5 Will Mn reduce Mg<sup>2+</sup> ions to Mg when all concentrations are 1.0M?

- 6.6 Will Cr<sub>2</sub>O<sub>7</sub><sup>2-</sup> oxidize Fe<sup>2+</sup> ions to Fe<sup>3+</sup> ions in acidic solution and under standard state conditions?

- 6.7 Which of the following reactions would occur spontaneously as written, when all substances are in their standard states?



- 6.8 Calculate  $E^\circ$  for the following reactions and predict whether the reactions should occur spontaneously as written when run under standard conditions:



- 6.9 (a) Will I<sub>2</sub> oxidize Fe<sup>2+</sup> to Fe<sup>3+</sup> in acid solution?

- (b) Will I<sub>2</sub> oxidize H<sub>2</sub>SO<sub>3</sub> to SO<sub>4</sub><sup>2-</sup> in acid solution?

$$E^\circ_{\text{H}_2\text{O}/\text{H}_2} = -0.828 \text{ V}$$

$$E^\circ_{\text{Fe}^{2+}/\text{Fe}^{3+}} = -0.770 \text{ V}$$

$$E^\circ_{\text{I}_2/\text{I}^-} = 0.535 \text{ V}$$

$$210 \quad E^\circ_{\text{Ca}/\text{Ca}^{2+}} = -2.76 \text{ V}$$

$$E^\circ_{\text{Fe}^{3+}/\text{Fe}^{2+}} = 0.77 \text{ V}$$

$$E^\circ_{\text{CrO}_4^{2-}/\text{Cr(OH)}_3} = +0.12 \text{ V}$$

$$E^\circ_{\text{Ni}/\text{Ni}^{2+}} = -0.72 \text{ V}$$

## Electrochemistry

- (c) Will  $\text{Sn}^{4+}$  oxidize  $\text{H}_2\text{O}_2$  to  $\text{O}_2$  in acid solution?  
 (d) Will  $\text{H}_2\text{O}$  reduce  $\text{Cl}_2$  to  $\text{Cl}^-$  in acid solution?
- Note that all reactions are run under standard conditions.
- 6.10 Calculate the volume of gas (at STP) that will collect at the cathode and anode when water is electrolyzed for 45.0 minutes with a 12.0-amp current.
- 6.11 With 10C of electric charge, how many grams of silver metal can be prepared? [Ar:Ag = 107.868].
- 6.12 Describe what happens during the electrolysis of the following aqueous solutions.  
 (a)  $\text{NaCl}$ ; (b)  $\text{Na}_2\text{SO}_4$ ; (c)  $\text{NaI}$ ; (d)  $\text{CaI}_2$  (e)  $\text{CuBr}_2$
- 6.13 Consider the following cathode reaction  

$$\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightarrow \text{Mn}^{2+} + 4\text{H}_2\text{O}$$
  
 Calculate the initial concentration of the permanganate solution if passage of 0.600 - amp current for 10 minutes completely reduced all the permanganate present in 15.0mL of solution.
- 6.14 Use the Nernst equation to calculate the cell potential at 25°C for the following cells:  
 (a)  $\text{Zn(s)}/\text{Zn}^{2+}(0.100\text{M})/\text{Ag}^+(0.010\text{M})/\text{Ag(s)}$        $E^\circ_{\text{Zn}^{2+}/\text{Zn}} = +0.763\text{V}$   
 (b)  $\text{Cu(s)}/\text{Cu}^{2+}(0.075\text{M})/\text{Ag}^+(0.015\text{M})/\text{Ag(s)}$        $E^\circ_{\text{Ag}^+/\text{Ag}} = +0.800\text{V}$
- 6.15 Calculate the cell potential for a voltaic cell composed of a piece of chromium metal immersed in a 0.240M solution of  $\text{Cr}^{3+}$  and a piece of zinc in a solution of 0.0240M of  $\text{Zn}^{2+}$ . Which electrode is the cathode?       $E^\circ_{\text{Zn}^{2+}/\text{Zn}} = -0.763\text{V}$        $E^\circ_{\text{Cr}^{3+}/\text{Cr}^{2+}} = +0.74\text{V}$
- 6.16 For a cell composed of  $\text{Cu}/\text{Cu}^{2+}$  electrode and  $\text{Fe}/\text{Fe}^{2+}$  electrode, the overall cell potential is 0.80V when the concentration of  $\text{Cu}^{2+}$  solution is 0.0500 M. What is the concentration of the  $\text{Fe}^{2+}$  solution?       $E^\circ_{\text{Cu}^{2+}/\text{Cu}} = -0.34\text{V}$        $E^\circ_{\text{Fe}^{2+}/\text{Fe}} = -0.41\text{V}$
- 6.17 Calculate the potential of the following cell when 99% of the zinc metal and  $\text{Cu}^{2+}$  ions have been consumed:  

$$\text{Zn/Zn}^{2+}(1.00\text{M})/\text{Cu}^{2+}(1.00\text{M})/\text{Cu}$$
- 6.18 Write the Nernst equation and determine the cell potential for the following reaction at 25°C.

## Electrochemistry

$$E^\circ_{Fe/Fe^{3+}} = +0.036\text{ V}$$

$$E^\circ_{Al^{3+}/Al} = -1.706\text{ V}$$

- 6.19  $Fe/Fe^{3+}(3.12M)/Al^{3+}(1.50M)/Al$   
 Ferrous ions are oxidised by silver ions according to the equation  
 $Fe^{2+}(aq) + Ag^+(aq) \rightleftharpoons Fe^{3+}(aq) + Ag(s)$
- Write a line notation for the cell
  - Write the half-cell reactions (that is, equations for the anode and cathode reactions).
  - Calculate the cell potential under standard-state conditions, and hence the standard free-energy change for the cell reaction, given that the equilibrium constant at 298 K is 7.810  
 $(F = 96500\text{ C mol}^{-1}, R = 8.314\text{ J mol}^{-1}\text{K}^{-1})$ .

- 6.20 Calculate the equilibrium constant at 25°C for the following reactions.

- $Cu^{2+}(aq) + Zn(s) \rightleftharpoons Cu(s) + Zn^{2+}(aq)$
- $Cu(s) + Br_2(aq) \rightleftharpoons Cu^{2+}(aq) + 2Br^-(aq)$
- $Zn(s) + 2H^+(aq) \rightleftharpoons Zn^{2+}(aq) + H_2(g)$
- $MnO_4^-(aq) + 5Cl^- + 8H^+ \rightleftharpoons Mn^{2+}(aq) + 5/2Cl_2(g) + 4H_2O(l)$
- $Cl_2(g) + 2Br^-(aq) \rightleftharpoons Br_2 + 2Cl^-(aq)$
- $Ag(s) + 2H^+(aq) \rightleftharpoons Ag^+(aq) + 1/2H_2(g)$
- $2Ag(s) + 2H^+(aq) \rightleftharpoons 2Ag^+(aq) + H_2(g)$

- 6.21 Calculate  $\Delta G^\circ$  from the standard state cell potential,  $E^\circ_{cell}$ , determined in question 16.20(a).

- 6.22 Calculate  $\Delta G^\circ$  for the following reaction at 25°C.



Will the reaction be spontaneous at this temperature?

- 6.23 Using the standard-state cell potentials given below, calculate the solubility products of (a)  $AgCl$ ; (b)  $Mg(OH)_2$ ; (c)  $Cu(OH)_2$ ; (d)  $Hg_2Cl_2$

