

SITE MENU *close*

SEARCH

Custom Search



NAVIGATION

- [Home](#)

SOCIAL

- [Facebook](#)
- [Twitter](#)
-

[up](#)



[Free Mathematics Tutorials](#)

Mathematical Induction - Problems With Solutions

Several problems with detailed solutions on mathematical induction are presented.

The principle of mathematical induction is used to prove that a given proposition (formula, equality, inequality...) is true for all positive integer numbers greater than or equal to some integer N.

Let us denote the proposition in question by $P(n)$, where n is a positive integer.

The proof involves two steps:

Step 1: We first establish that the proposition $P(n)$ is true for the lowest possible value of the positive integer n .

Step 2: We assume that $P(k)$ is true and establish that $P(k+1)$ is also true

Problem 1

Use mathematical induction to prove that

$$1 + 2 + 3 + \dots + n = n(n + 1) / 2$$

for all positive integers n .

Solution to Problem 1:

Let the statement $P(n)$ be

$$1 + 2 + 3 + \dots + n = n(n + 1) / 2$$

STEP 1: We first show that $p(1)$ is true.

Left Side = 1

Right Side = $1(1 + 1) / 2 = 1$

Both sides of the statement are equal hence $p(1)$ is true.

STEP 2: We now assume that $p(k)$ is true

$$1 + 2 + 3 + \dots + k = k(k + 1) / 2$$

and show that $p(k + 1)$ is true by adding $k + 1$ to both sides of the above statement

$$\begin{aligned} 1 + 2 + 3 + \dots + k + (k + 1) &= k(k + 1) / 2 + (k + 1) \\ &= (k + 1)(k / 2 + 1) \\ &= (k + 1)(k + 2) / 2 \end{aligned}$$

The last statement may be written as

$$1 + 2 + 3 + \dots + k + (k + 1) = (k + 1)(k + 2) / 2$$

Which is the statement $p(k + 1)$.

Problem 2

Prove that

$$1^2 + 2^2 + 3^2 + \dots + n^2 = n(n + 1)(2n + 1)/6$$

For all positive integers n.

Solution to Problem 2:

Statement P (n) is defined by

$$1^2 + 2^2 + 3^2 + \dots + n^2 = n(n + 1)(2n + 1)/2$$

STEP 1: We first show that $p(1)$ is true.

$$\text{Left Side} = 1^2 = 1$$

$$\text{Right Side} = 1(1 + 1)(2*1 + 1)/6 = 1$$

Both sides of the statement are equal hence $p(1)$ is true.

STEP 2: We now assume that $p(k)$ is true

$$1^2 + 2^2 + 3^2 + \dots + k^2 = k(k + 1)(2k + 1)/6$$

and show that $p(k + 1)$ is true by adding $(k + 1)^2$ to both sides of the above statement

$$1^2 + 2^2 + 3^2 + \dots + k^2 + (k + 1)^2 = k(k + 1)(2k + 1)/6 + (k + 1)^2$$

Set common denominator and factor $k + 1$ on the right side

$$= (k + 1)[k(2k + 1) + 6(k + 1)]/6$$

Expand $k(2k + 1) + 6(k + 1)$

$$= (k + 1) [2k^2 + 7k + 6] / 6$$

Now factor $2k^2 + 7k + 6$.

$$= (k + 1) [(k + 2)(2k + 3)] / 6$$

We have started from the statement $P(k)$ and have shown that

$$1^2 + 2^2 + 3^2 + \dots + k^2 + (k + 1)^2 = (k + 1) [(k + 2)(2k + 3)] / 6$$

Which is the statement $P(k + 1)$.

Problem 3

Use mathematical induction to prove that

$$1^3 + 2^3 + 3^3 + \dots + n^3 = n^2(n + 1)^2 / 4$$

for all positive integers n .

Solution to Problem 3:

Statement $P(n)$ is defined by

$$1^3 + 2^3 + 3^3 + \dots + n^3$$

$$+ \dots + n^3 = n^2(n+1)^2 / 4$$

STEP 1: We first show that $p(1)$ is true.

$$\text{Left Side} = 1^3 = 1$$

$$\text{Right Side} = 1^2(1+1)^2 / 4 = 1$$

hence $p(1)$ is true.

STEP 2: We now assume that $p(k)$ is true

$$1^3 + 2^3 + 3^3 + \dots + k^3 = k^2(k+1)^2 / 4$$

add $(k+1)^3$ to both sides

$$1^3 + 2^3 + 3^3 + \dots + k^3 + (k+1)^3 = k^2(k+1)^2 / 4 + (k+1)^3$$

factor $(k+1)^2$ on the right side

$$= (k + 1)^2 [k^2 / 4 + (k + 1)]$$

set to common denominator and group

$$= (k + 1)^2 [k^2 + 4k + 4] / 4$$

$$= (k + 1)^2 [(k + 2)^2] / 4$$

We have started from the statement $P(k)$ and have shown that

$$1^3 + 2^3 + 3^3 + \dots + k^3 + (k + 1)^3 = (k + 1)^2 [(k + 2)^2] / 4$$

Which is the statement $P(k + 1)$.

Problem 4

Prove that for any positive integer number n , $n^3 + 2n$ is divisible by 3

Solution to Problem 4:

Statement $P(n)$ is defined by _____

$n^3 + 2n$ is divisible by 3

STEP 1: We first show that $p(1)$ is true. Let $n = 1$ and calculate $n^3 + 2n$

$$1^3 + 2(1) = 3$$

3 is divisible by 3

hence $p(1)$ is true.

STEP 2: We now assume that $p(k)$ is true

$k^3 + 2k$ is divisible by 3

is equivalent to

$k^3 + 2k = 3M$, where M is a positive integer.

We now consider the algebraic expression $(k+1)^3 + 2(k+1)$; expand it and group like terms

$$(k+1)^3 + 2(k+1) = k^3 + 3k^2 + 5k + 3$$

$$= [k^3 + 2k] + [3k^2 + 3k + 3]$$

$$= 3M + 3[k^2 + k + 1] = 3[M + k^2 + k + 1]$$

Hence $(k+1)^3 + 2(k+1)$ is also divisible by 3 and therefore statement $P(k+1)$ is true.

Problem 5

Prove that $3^n > n^2$ for $n = 1, n = 2$ and use the mathematical induction to prove

that $3^n > n^2$ for n a positive integer greater than 2.

Solution to Problem 5:

Statement $P(n)$ is defined by

$$3^n > n^2$$

STEP 1: We first show that $p(1)$ is true. Let $n = 1$ and calculate 3^1 and 1^2 and compare them

$$3^1 = 3$$

$$1^2 = 1$$

3 is greater than 1 and hence $P(1)$ is true.

Let us also show that $P(2)$ is true.

$$3^2 = 9$$

$$2^2 = 4$$

Hence $P(2)$ is also true.

STEP 2: We now assume that $P(k)$ is true

$$3^k > k^2$$

Multiply both sides of the above inequality by 3

$$3 * 3^k > 3 * k^2$$

The left side is equal to 3^{k+1} . For $k > 2$, we can write

$$k^2 > 2k \text{ and } k^2 > 1$$

We now combine the above inequalities by adding the left hand sides and the right hand sides of the two inequalities

$$2k^2 > 2k + 1$$

We now add k^2 to both sides of the above inequality to obtain the inequality

$$3k^2 > k^2 + 2k + 1$$

Factor the right side we can write

$$3 * k^2 > (k + 1)^2$$

If $3 * 3^k > 3 * k^2$ and $3 * k^2 > (k + 1)^2$ then

$$3 * 3^k > (k + 1)^2$$

Rewrite the left side as 3^{k+1}

$$3^{k+1} > (k+1)^2$$

Which proves that $P(k+1)$ is true

Problem 6

Prove that $n! > 2^n$ for n a positive integer greater than or equal to 4. (Note: $n!$ is n factorial and is given by $1 * 2 * \dots * (n-1)*n$.)

Solution to Problem 6:

Statement $P(n)$ is defined by

$$n! > 2^n$$

STEP 1: We first show that $p(4)$ is true. Let $n = 4$ and calculate $4!$ and 2^4 and compare them

$$4! = 24$$

$$2^4 = 16$$

24 is greater than 16 and hence $p(4)$ is true.

STEP 2: We now assume that $p(k)$ is true

$$k! > 2^k$$

Multiply both sides of the above inequality by $k+1$

$$k!(k+1) > 2^k(k+1)$$

The left side is equal to $(k+1)!$. For $k > 4$, we can write

$$k+1 > 2$$

Multiply both sides of the above inequality by 2^k to obtain

$$2^k(k+1) > 2 * 2^k$$

The above inequality may be written

$$2^k(k+1) > 2^{k+1}$$

We have proved that $(k+1)! > 2^k(k+1)$ and $2^k(k+1) > 2^{k+1}$ we can now write

$$(k+1)! > 2^{k+1}$$

We have assumed that statement $P(k)$ is true and proved that statement $P(k+1)$ is also true.

Problem 7

Use mathematical induction to prove De Moivre's theorem

$$[R(\cos t + i \sin t)]^n = R^n(\cos nt + i \sin nt)$$

for n a positive integer.

Solution to Problem 7:

STEP 1: For $n = 1$

$$[R(\cos t + i \sin t)]^1 = R^1(\cos 1*t + i \sin 1*t)$$

It can easily be seen that the two sides are equal.

STEP 2: We now assume that the theorem is true for $n = k$, hence

$$[R(\cos t + i \sin t)]^k = R^k(\cos kt + i \sin kt)$$

Multiply both sides of the above equation by $R(\cos t + i \sin t)$

$$[R(\cos t + i \sin t)]^k R(\cos t + i \sin t) = R^k (\cos kt + i \sin kt) R(\cos t + i \sin t)$$

Rewrite the above as follows

$$[R(\cos t + i \sin t)]^{k+1} = R^{k+1} [(\cos kt \cos t - \sin kt \sin t) + i(\sin kt \cos t + \cos kt \sin t)]$$

Trigonometric identities can be used to write the trigonometric expressions $(\cos kt \cos t - \sin kt \sin t)$ and $(\sin kt \cos t + \cos kt \sin t)$ as follows

$$(\cos kt \cos t - \sin kt \sin t) = \cos(kt + t) = \cos(k + 1)t$$

$$(\sin kt \cos t + \cos kt \sin t) = \sin(kt + t) = \sin(k + 1)t$$

Substitute the above into the last equation to obtain

$$[R(\cos t + i \sin t)]^{k+1} = R^{k+1} [\cos(k + 1)t + \sin(k + 1)t]$$

It has been established that the theorem is true for $n = 1$ and that if it assumed true for $n = k$ it is true for $n = k + 1$.

More References and links

[maths problems with detailed solutions](#).

[Home Page](#)



report this ad

Popular Pages

- [Math Problems, Questions and Online Self Tests](#)
- [File Not Found](#)
- [Geometric Sequences Problems with Solutions](#)
- [Questions on Functions with Solutions](#)
- [Arithmetic Sequences Problems with Solutions](#)

[back to top](#)

• [Privacy Policy](#)