MTH 103

Tutorial

Questions

Temiwhite XYB

1. The tangent of the parabola y^2 =4ax at the point (at ² ,2at) meets the x-axis at T. The straight line through P parallel to the axis of the parabola meets the directrix at Q. If M is the mid-point of PT and N is the mid-point of PM. Find the equation of the locus of M
a. 3 b. 2 c. 0 d. 1
2. The tangent of the parabola y^2 =4ax at the point (at²,2at) meets the x-axis at T. The straight line through P parallel to the axis of the parabola meets the directrix at Q. If M is the mid-point of PT and N is the mid-point of PM. Find the equation of the locus of N
a. $2y^2=9ax^2$ b. $-2y^2=9ax$ c. $2y^2=-9ax$ d. $2y^2=9ax$
3. Find the equations of the tangents and the normal at the point (4,1) on the curve xy=4
a4y+x=8, y-4x+15=0 b. 4y+x=8, y-4x-15=0c. 4y-x=8, y-4x+15=0d. 4y+x=8, y-4x+15=0
4. Obtain the equation of the normal to the parabola $y^2=4ax$ at the point $(at^2,2at)$
a. $-y^2=a(x-3a)$ b. $y^2=a(x+3a)$ c. $y^2=a(x-3a)$ d. $y^2=-a(x-3a)$
5. If the parabola y^2 =4ax cuts the hyperbola $xy=c^2$ at right angles, find the coordinate of the point of intersection.
a. (-2a, -2Sqrt(2a)) b. (2a, 2Sqrt(2a)) c. (2a, 2Sqrt(-2a)) d. (-2a, 2Sqrt(2a))
6. If circles $x^2+y^2+2x-8y+8=0$ and $x^2+y^2+10x-2y+22=0$ touhes one another, find the point of contact
a. (-17/5,11/5) b. (-17/5,-11/5) c. (17/5,11/5) d. (-17,11)
7. If circles $x^2+y^2+2x-8y+8=0$ and $x^2+y^2+10x-2y+22=0$ touhes one another, find the equation to the common tangent at this point
a. 4x+3y+7=0 b. 4x-3y+7=0 c. 4x+3y-7=0 d. 4x-3y-7=0
8. If circles $x^2+y^2+2x-8y+8=0$ and $x^2+y^2+10x-2y+22=0$ touhes one another, find the area of the triangle enclosed by this common tangent, the line of centres and the y-axis
a. 12 square units b. $12^{1}/_{24}$ square units c. 12.5 square units d. $-12^{1}/_{24}$ square units
9. The equation of the tangents to the hyperbola $x^2-y^2=7$ which are parallel to $3y=4x$. Find their points of contact
a. (3,3) b. (4,4) c. (4,3) d. (3,4)
10. The equation of the tangents to the hyperbola $x^2-y^2=7$ which are parallel to $3y=4x$. Find the
area of the triangle which one of these tangents makes with the asymptotes.


b7 square un	its c. 6 square	units d. 7 squ	are units
e from a fixed str	aight line ZQ (the	directrix) is a co	nstant (e, called
. Hyperbola (c). Parabola	(d). Elli	pse
e from a fixed str	raight line ZQ (the	e directrix) is a co	onstant (e, called
71			
ce from a fixed st	raight line ZQ (th	e directrix) is a c	onstant (e, called
o). Parabola (c). Elilipse	(d). Hy	perbola
of the tangents to	the parabola $y^2=4$	8x at the points (3,12), (48,-48).
(b). $y=x/2-24$	(c). $y=-x/2-24$	(d). $y=-x/2+24$	
² =48x at the poin	ts (3,12), (48,-48)	. Find their point	of intersection .
8 (b). a=12, x=-1	2, y=-18 (c). a=1	.2, x=-12, y=18	(d). a=12, x=12, y=18
s of the tangents f	from the point (2,4	4) to the parabola	$y^2=6x$ at m=3/2.
		Α-	
3/5 (c). 5/3		5/3	
ates of the foci to	the ellipse $x^2/25$ +	$y^2/16=1$	
(b). (-3,0)			
	P(x,y) which more from a fixed strent forms according the Hyperbola (at P(x,y) which more from a fixed strent forms according to the tangents to (b). Parabola (c) Parabola (d) Parabola (e) Parabola (P(x,y) which moves so that the rate of from a fixed straight line ZQ (the ent forms according to the value of the P(x,y) which moves so that the rate of from a fixed straight line ZQ (the ent forms according to the value of the P(x,y) which moves so that the rate of from a fixed straight line ZQ (the ent forms according to the value of the P(x,y) which moves so that the rate of from a fixed straight line ZQ (the ent forms according to the value of the tangents to the parabola $y^2 = 4$ (b). $y = x/2 - 24$ (c). $y = -x/2 - 24$ (d). $y = x/2 - 24$ (e). $y = -18$ (for the tangents from the point (2,4). $y = x/2 - 3$ (c). $y = 3x/2 + 3$ (d) as of the tangents from the point (2,4). $y = x/2 - 3$ (c). $y = 3x/2 + 3$ (d) at the foci to the ellipse $x^2/25 + y^2/16 = 1$ (d) at the foci to the ellipse $x^2/25 + y^2/16 = 1$ (d) at the foci to the ellipse $x^2/25 + y^2/16 = 1$ (d) at the foci to the ellipse $x^2/25 + y^2/16 = 1$ (d) at the foci to the ellipse $x^2/25 + y^2/16 = 1$ (d) at the foci to the ellipse $x^2/25 + y^2/16 = 1$ (d) at the foci to the ellipse $x^2/25 + y^2/16 = 1$ (d) at the foci to the ellipse $x^2/25 + y^2/16 = 1$ (d) at the foci to the ellipse $x^2/25 + y^2/16 = 1$ (d) at the foci to the ellipse $x^2/25 + y^2/16 = 1$ (d) at the foci to the ellipse $x^2/25 + y^2/16 = 1$ (d) at the foci to the ellipse $x^2/25 + y^2/16 = 1$ (d) at the foci to the ellipse $x^2/25 + y^2/16 = 1$ (d) at the foci to the ellipse $x^2/25 + y^2/16 = 1$ (e) at $x = 1$ (for $x = 1$) and $x =$	P(x,y) which moves so that the ratio of its distance of from a fixed straight line ZQ (the directrix) is a content forms according to the value of e. The locus is known that the ratio of its distance of the form a fixed straight line ZQ (the directrix) is a content form a fixed straight line ZQ (the directrix) is a content forms according to the value of e. The locus is known to the value of e. The locus is known to the parabola (c). Elilipse (d). Paratrophysical transfer forms according to the value of e. The locus is known to the parabola (c). Elilipse (d). Hy of the tangents to the parabola y²=48x at the points (d). Hy of the tangents to the parabola y²=48x at the points (d). y=x/2-24 (e). y=-x/2-24 (d). y=-x/2+24 (e). y=-x/2-24 (d). y=-x/2+24 (e). a=12, x=-12, y=18 (e). a=12, x=-12, y=18 (e). a=12, x=-12, y=18 (e). a=12, x=-12, y=18 (f). y=x/2-3 (f). y=3x/2+3 (f). y=3x/2+1 (f). y=x/2-3 (f). y=3x/2+3 (f). y=x/2-3 (f). y=3x/2+3 (f). y=x/2-3 (f). y=3x/2+3 (f). y=x/2-3 (f). y=x/

Ografia 1 + los 1 ig (c). (+25/3) (d). (+ 25/3)

20. Find the equation of the directrices of the ellipse $x^2/25+y^2/16=1$

(b). (-25/3)

(a). (3/25)



30. Find the equation of the normal to the parabola y^2 =4ax at the point (at ² ,2at). P,Q are two points on the parabola such that the chord PQ Subtends a right angle at the vertex of the parabola. Find the locus of the point of intersection of the normals at P and Q.				
a. $y^2=-16a(x-2a)$ b. $y^2=16a(x+2a)$ c. $y^2=-16a(x+2a)$ d. $y^2=16a(x-2a)$				
31. The asymptotes will each be inclined at to the x-axis.				
(a). 90^0 (b). 45^0 (c). 180^0 (d). 60^0				
32. Find the centre and radius of $x^2+y^2+2x+6y+6=0$				
a. C(-1,-3), r=2 b. C(-1,3), r=2 c. C(1,-3), r=2 d. C(1,-3), r=2				
33. Find the equation of the circle center $(7,-6)$ which touches the line $3x-4y+5=0$				
(a) $x^2+y^2-14x+12y-15=0$. (b) $x^2-y^2-14x+12y-15=0$. (c) $x^2+y^2-14x+12y+15=0$. (d) $x^2+y^2+14x+12y-15=0$.				
34. Find the equation of the circle passing through the points (0,0), (3,1), (5,5)				
a. $x^2-y^2+10y=0$ b. $x^2+y^2-10y=0$ c. $x^2-y^2-10y=0$ d. $x^2+y^2+10y=0$				
35. Find the equation of the tangent to the circle $2x^2+2y^2+x-11y=0$ at (-2,5)				
a. $7x+9y-59=0$ b. $7x+9y+59=0$ c. $7x-9y+59=0$ d. $7x-9y-59=0$				
36. Find the point of intersection of the line $x+y-3=0$ and the circle $x^2+y^2+x-5y+4=0$				
a. (-1,2), (-1,4) b. (1,-2), (-1,4) c. (1,2), (-1,-4) d. (1,2), (-1,4)				
37. The length of the tangent from the point (3,2) to the circle $x^2+y^2-2x-3y+k=0$ is 9units. Find he value of k.				
a. 88 b. 81 c. 8 d. 80				
38. Given the three circles $x^2+y^2-16x+60=0$, $x^2+y^2-12x+20=0$ and $x^2+y^2-16x-12y+84=0$. Find the length of each tangent.				
a. 4units b. 1unit c. 2units d. 3units				
39. Determine the two values of c for which the line $3x+4y+c=0$ is a tangent to the circle $x^2+y^2-6x-2y-15=0$				
a12 or 38 b. 12 or -38 c12 or -38 d. 12 or 38				
40. Give Circles $x^2+y^2-20x-14y+113=0$ and $4x^2+4y^2+16x-16y-49=0$. Find the length of the shortest distance from a point on one circle to a point on the other.				

(m) an